

Dynamics of Mercury in Eared Grebes on the Great Salt Lake

Presentation to the Mercury Workgroup
November 15, 2007

Nathan Darnall, USFWS
Keith Miles, USGS BRD
Dr. Joe Jehl, Smithsonian



Funded Study FY'06

- U.S. Fish and Wildlife, Ecological Services (UT)
- U.S. Geological Survey, Biological Resources Division, U.C. Davis (CA)
- Dr. Joe R. Jehl, Smithsonian Institute


Partners

- U.S. Geological Survey, WRD (UT) (WI)
- Utah Division of Water Quality
- Utah Division of Wildlife Resources

Eared Grebe (C. Allan Morgan)



Distribution of the Eared Grebe in North and Central America.



Why *Podiceps nigricollis*?

- Eared grebes spend 3 to 4 months on GSL during fall migration (Sept – Dec)
- They eat brine shrimp almost exclusively
- Previous FWS data suggest Hg accumulation
- There is a reference population at Mono Lake in California
- Historic data and archived samples

Study Questions

- Does mercury accumulate in grebes residing on Great Salt Lake?
- If so, is GSL different than other lakes?
- Does molting provide a way for grebes to eliminate excess mercury?
- Have conditions at GSL changed over time?

Additional Considerations

- Are other species that utilize GSL at risk?
- Are there implications for human health?
- What role does molting and feather growth play in other species?
- Where might the GSL be headed?


Hypotheses

- Do Hg concentrations differ:
 - 1) Among stage of migration (Oct / Dec)
 - 2) Among locations (GSL / Mono Lake)
 - 3) Between years (1992 / 2006)


Collections

- Water (unfiltered, total and methyl)
- Brine Shrimp (total)
- Eared Grebes
 - Livers (total and methyl)
 - Breast Muscle (total and methyl)
 - Breast Feathers (total)
 - Primary Feathers (total)

How to Read a Box Plot








Mercury in Unfiltered Great Salt Lake Water






Trolling for brine shrimp.



Brine shrimp streak.





Conclusions

- Mercury increased in livers, breast muscle and breast feathers of birds on GSL but only livers were significant (<0.05)
- Concentrations were significantly higher at GSL compared to Mono Lake for all tissues
- Concentrations in livers were significantly higher in 2006 compared to 1992

Conclusions

- Total and methyl mercury in unfiltered water samples were significantly higher in December samples
- Mercury in brine shrimp was significantly higher in December compared to October
- Concentrations in brine shrimp were three times higher than a decade ago

